REPORT DOCUMENTATION PAGE

Unclassified Unclassified Unclassified SAR

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204 Actions NA 22202-4302 and to the Office of Management and Budget Paperwork Reduction Project (0704-0188) Washington, DC 20503.

			agement and Budget, Paperwood. HE ABOVE ADDRESS.	ork Reduction Pro	ject (0704-0188),	Washington, DC 20503.		
1. REPORT DAT	E (DD-MM-YYYY)	2. RE	PORT TYPE	-		3. DATES COVERED		
			Conference	Proceeding		08-12 September 2003		
4. TITLE AND S	UBTITLE				5a. CON	ITRACT NUMBER		
About Nonlinear Dependence of Remote Sensing and Diffuse Reflection Coefficients on Gordon's Parameter				5b. GRA	5b. GRANT NUMBER			
			5c. PRO	5c. PROGRAM ELEMENT NUMBER				
6. AUTHOR(S)					5d. PRO	JECT NUMBER		
o. Ao mente,						VSF 73-6641-02-5		
V. I. Haltrin a	and Sonia C. G	allegos			5e. TAS	5e. TASK NUMBER		
			5f. WOR	5f. WORK UNIT NUMBER				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004			_	8. PERFORMING ORGANIZATION REPORT NUMBER				
•						NRL/PP/7330-03-0010		
		GENCY NAME(S)	AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
Office of Nav 800 N. Quinc						ONR		
Arlington, VA	Å 22217-5660					11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTI	ON/AILABILITY ST	TATEMENT						
Approved for	public release	; distribution is	s unlimited		2	0074947 900		
13. SUPPLEMENTARY NOTES 20031217 280								
14. ABSTRACT								
Remote sensing i~ and diffuse reflection R coefficients of seawater are dependent on inherent optical properties of seawater through the Gordon's parameter bB / (a bB), where a is an absorption and b is backscattering coefficients. g =+B Majority of researchers in ocean optics use linear approximation for ~ and R, i. e. ~g and R oc g. This approach works well when g ~ 0.1. All open and largest part of coastal waters satisfy this condition, but there are some cases when parameter g is as large as 0.98. We illustrate this fact with a histogram of Gordon's parameter g for the Yellow sea waters. We present a choice of alternative rigorous nonlinear equations for R that solve this problem, i.e. significantly reduce the error of the restoration of the Gordon's parameter g from remote sensing optical data and estimate possible errors in using linear dependencies								
15. SUBJECT TERMS nonlinear, remote sensing, diffuse reflection, Gordon's parameter								
16. SECURITY C	CLASSIFICATION	OF:	17. LIMITATION OF	18. NUMBER	19a. NAME OF	RESPONSIBLE PERSON		
a. REPORT	REPORT b. ABSTRACT c. THIS PAGE ABSTRACT OF Vladimir Haltrin PAGES 19h TELEPHONE NUMBER (lockude sees code)							

228-688-4528

PUBLICATION OR PRESE	NTATION RELEASE REQUEST		Pubkey: 3717 NRLINST 5600
REFERENCES AND ENGLOSURE	S Z TYPE OF HEBLICATION OF RESERVE		4 Anger Str. appeals a dictaling
Ref: (a) NRL Instruction 5600.2 (b) NRL Instruction 5510.40D Encl: (1) Two copies of subject pape (or abstract	() Book () Conference Proceedings (X) Conference () Conference () Conference () Multiple () Journal article (refereed)	ract only, not published a chapter rerence Proceedings trefereed) media report nal article (not refereed) Presentation, not publisl	hed
	() Other, explain	•	approval obtained X yesno
4. AUTHOR			
Title of Paper or Presentation			
Author(s) Name(s) (First, MI, Last), (mote Sensing and Diffuse Reflection Coefficie	nts on Gordon's Paran	neter
Vladimir I. Haltrin Sonia C. Galle			
t is intended to offer this paper to	the 2nd Int. ONW Conference	lome of Conformaci	
)8-SEP - 12-SEP-2003, St. Peters	·	ame of Conference)	
	(Date, Place and Classification o	f Conference)	
and/or for publication in 2nd Int.	ONW Conference, Unclassified		
· (/	lame and Classification of Publication)		(Name of Publisher)
with reference (a).	ertinent publication/presentation data will be	e entered in the public	cations data base, in accordance
This paper does not violate any dis	he subject paper (is $__$) (is not $_$) classclosure of trade secrets or suggestions of c	utside individuals or o	concerns which have been
This subject paper (has) (ha	n confidence. This paper (does) (does s never _X) been incorporated in an office	s not <u>X</u>) contain a ial NRL Report.	ny militarily critical technology.
		· \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Jallie.
Name and	imir I. Haltrin, 7333 d Code (<i>Principal Author</i>)	γνι	(Signature)
			And the second s
CODE Author(s)	SIGNATURE	DATE	COMMENTS
Haltrin	Mtaltriy	7/16/03	abstract to this paper is
			Uresdy Symen
Section Head	0 1 1 21 00	1/12	
Gould, acting Branch Head	w tould	7 16 03	
Arnone	Kleman	7/15/02	
Division Head		// /	Release of this paper is approved. To the best knowledge of this Division, the
Payne, acting	Sulph	7/17/03	subject matter of this paper (has) (has neverX) been classified.
Security, Code 7030.1	And Star la	18/07	1. Paper or abstract was released. 2. A copy is filed in this office 652° - 281 - 3
Office of Counsel,Code	The state of the s	1/10/03	53.701-3
ADOR/Director NCST E.O. Hartwig, 7000	tuesty fait	1/2/103	
ublic Affairs (Unclassified/ Inlimited Only), Code 7030.4			
ivision, Code			
uthor, Code			

U, L	NOT THE THE PROPERTY OF THE PR	The second secon					
A - Approved for public release, distribution is unlimited.							
	B - Distribution authorized to U.S. Government						
	Foreign Government Information	Contractor Performance Evaluation	Critical Technology				
	Proprietary Information	Administrative/Operational Use	Premature Dissemination				
	Test and Evaluation	Software Documentation	Cite "Specific Authority"				
	Date statement applied		(Identification of valid documented authority)				
İ	Other requests for this document shall be	e referred to(Insert Controlling DOI	O Office*)				
	C - Distribution authorized to U.S. Governmen	-	eason below):				
	Foreign Government Information	Software Documentation					
l	Administrative/Operational Use	Critical Technology	Cite "Specific Authority (Identification of valid documented authority)				
	Date statement applied						
	Other requests for this document shall be		D.Office #1				
l		(Insert Controlling DOI	of Office)				
	D - Distribution authorized to DOD and DOD c						
	Foreign Government Information	Critical Technology	n				
	Software Documentation	Cite "Specific Authority	of valid documented authority)				
	Administrative/Operational Use	(,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	···				
	Date statement applied	<u></u>					
	Other requests for this document shall b	e referred to	0.05.4				
١.		(Insert Controlling DO	D Office")				
	E - Distribution authorized to DOD componen		Critical Technology				
	Proprietary Information	Premature Dissemination					
ŀ	Foreign Government Information	Software Documentation	Direct Military Support				
	Administrative/Operational Use	Contractor Performance Evaluation	Test and Evaluation				
	Date statement applied	-	Cite "Specific Authority" (Identification of valid documented authority)				
l	Other requests for this document shall t	e referred to	(Identification of Valid documented database)				
1		(Insert Controlling DO	D Office*)				
	F - Further dissemination only as directed by						
1		(Insert Controlling	DOD Office")				
	Date statement applied	or higher DOD authority					
	G - Distribution authorized to U.S. Governme	nt agencies and private individuals or ent	erprises eligible to obtain export-controlled				
1	technical data in accordance with regulati	ions implementing 10 U.S.C. 140c.					
1	Date statement applied						
	Other requests for this document shall be referred to						
1	(Insert Controlling DOD Office*) *For NRL publications, this is usually the Commanding Officer, Naval Research Laboratory, Washington, DC 20375-5320						
	OTHER LIMITATION AT LINES LE MILITARIA		i Geolificali merstemi i Relegii sadiktaring selelik kalan bili ilgi malik seleti.				
	Classificat	ion NOFORN	DTIC exempt (explain)				
1.	Chariffeetien Baylow Substantive ch	anges made in this document after appro	val by Classification Review and Public Release invalidate				
	Classification Review	Therefore, if any substantive changes are nust be returned for another Classificatio	made by the author, Technical Information, or anyone else,				
L	and documents						
8	NSTRUCTORS FOR THE STATE OF THE						
Г			vision head for review and approval according to				
	the routing in Section 4. 1. NRL Reports	Submit ti	ne diskette (if available), manuscript, typed double-spaced, with tables, illustrations, references, draft SF 298, and				
1		nronosed	distribution list.				
	2. NRL Memorandum ReportsSubmit a copy of the original, typed manuscript complete with tables, illustrations, references, draft SF 298, and proposed						
	AIDI Dublination or allow heater has	distribution					
	proceedings, or any other printed pul	blications.	·				

ABOUT NONLINEAR DEPENDENCE OF REMOTE SENSING AND DIFFUSE REFLECTION COEFFICIENTS ON GORDON'S PARAMETER

Vladimir I. Haltrin, and Sonia C. Gallegos

Naval Research Laboratory, Ocean Optics Section, Code 7333, Stennis Space Center, MS 39529-5004, USA e-mail: haltrin@nrlssc.navy.mil, web page: http://haltrin.freeshell.org

Abstract: Remote sensing r_{rs} and diffuse reflection R coefficients of seawater are dependent on inherent optical properties of seawater through the Gordon's parameter $g = b_B / (a + b_B)$, where a is an absorption, and b_B is backscattering coefficients. Majority of researchers in ocean optics use linear approximation for r_{rs} and R, i. e. $r_{rs} \propto g$ and $R \propto g$. This approach works well when $g \leq 0.1$. All open and largest part of coastal waters satisfy this condition, but there are some cases when parameter g is as large as 0.98. We illustrate this fact with a histogram of Gordon's parameter g for the Yellow sea waters. We present a choice of alternative rigorous nonlinear equations for R that solve this problem, i.e. significantly reduce the error of the restoration of the Gordon's parameter g from remote sensing optical data and estimate possible errors in using linear dependencies

1. Introduction

Diffuse reflection coefficient (DRC) of water body is an informative part of remote sensing reflectance [1] of light by the ocean. DRC contains information on content of dissolved and suspended substances in seawater. DRC is an apparent optical property that depends not only on inherent optical properties of the seawater, but also on the parameters of illumination. The dependence on inherent optical properties is expressed through the dependence on Gordon's parameter, i.e. the ratio of backscattering coefficient b_B to the sum of absorption a and backscattering coefficient, $g = b_B / (a + b_B)$. In the open ocean DRC is linearly proportional to g. This linear equation is very good for the Type I open ocean waters [2]. It is also acceptable for about 90% of coastal waters. Theoretical and numerical analysis show that the linear relationship can be adequately used only when Gordon's parameter g is relatively small, i.e. g < 0.1. This criterion is always satisfied in open ocean waters. The available database of experimental measurements show that in coastal waters Gordon's parameter may exceed this critical value of 0.1. In some very turbid coastal waters it can even reach values higher than 0.95. For example, in waters of Yellow Sea or coastal waters close to river estuaries the percentage of cases when g > 0.1 can reach 50% or more. In this paper we illustrate this fact with a histogram of Gordon's parameter for the Yellow sea waters. We estimate possible errors in using the linear dependence and present a choice of alternative rigorous nonlinear equations for DRC that solve this problem, i.e. significantly reduce the error of the restoration of the Gordon's parameter from remote sensing optical data.




Figure 1. Distribution of occurrences of measured Gordon's parameter in Yellow Sea in 2000.

2. Gordon's parameter in highly scattering marine environment

Until recently we had no significant cases of *in situ* measurements that show existence of sea waters with high cases of Gordon's parameter g exceeding 0.1. Relatively recent measurements made in 2000 by NRL researches with collaboration of Korean scientists in Yellow Sea shows that these highly scattering waters exhibit these characteristics. Figure 1 shows a histogram of the frequency of measurements of Gordon's parameter in this expedition. In this case more than 50% of measurements correspond to Gordon's parameter exceeding critical limit of linear approach. It means that processing of such data should involve nonlinear equations connecting DRC with g.

3. Equations for diffuse reflectance that valid for all values of Gordon's parameter

Fortunately, we have both experimental data and theoretical equations that correctly connect diffuse reflection coefficient with Gordon's parameter g, and, consequently, with a and b_B for the full range of their variability, $0 \le g \le 1$, $0 \le a \le \infty$, $0 \le b_B \le \infty$. First, we have experimental data published by Timofeyeva in 1972 and 1979 [3, 4] (see Tab. 1). Second, we have several equations that connect DRC with g in the full range of variability of this parameter, $0 \le g \le 1$. Let us consider these cases.

Table 1. Optical properties of natural and modeled scattering and absorbing media according to experiments by Timofeyeva. Here $\overline{\mu}$, $\overline{\mu}_d$, and $\overline{\mu}_u$ are, correspondingly, total, downward, and upward average cosines over radiance distribution in the scattering medium [5].

$oxedsymbol{\overline{\mu}}$	$\overline{\mu}_{\!\scriptscriptstyle d}$	$\overline{\mu}_{\!\scriptscriptstyle u}$	R	g	R/g
0	0.5	0.5	1.0	1.0	1.0
0.1	0.5249	0.4831	0.671	0.9408	0.7132
0.2	0.5525	0.4545	0.443	0.7970	0.5550
0.3	0.5834	0.4202	0.283	0.6179	0.4580
0.4	0.6184	0.3745	0.171	0.4439	0.3852
0.5	0.6566	0.3311	0.095	0.2959	0.3211
0.6	0.7008	0.3003	0.048	0.1802	0.2664
0.7	0.7536	0.2857	0.0207	0.0967	0.2141
0.8	0.8217	0.3610	0.0082	0.0413	0.1985
0.9	0.9033	0.6849	0.0016	0.0101	0.1584
1.0	1.0	1.0	0.0	0.0	0.25

a) Equation derived from the exact solution of radiative transfer equation in the depth of scattering medium [6]:

$$R = \frac{1 - \eta}{1 + \eta} \left(\sqrt{1 + \eta^2} - \eta \right)^2, \quad \eta = \sqrt{\frac{1 - g}{1 + (3 + 2\sqrt{2})g}} = \sqrt{\frac{a}{a + 2(2 + \sqrt{2})b_R}}.$$
 (1)

b) Equation derived in the framework of self-consistent approach [5] (asymptotic case):

$$R_{\infty} = \left(\frac{1 - \overline{\mu}}{1 + \overline{\mu}}\right)^2,\tag{2}$$

$$\overline{\mu} = \sqrt{\frac{1 + 2g - \sqrt{g(4 + 5g)}}{1 + g}} = \sqrt{\frac{a}{a + 3b_B + \sqrt{b_B(4a + 9b_B)}}}.$$
 (3)

c) Equation derived in the framework of self-consistent approach [5] (case of diffuse illumination):

Proceedings of the II International Conference "Current Problems in Optics of Natural Waters," ONW'2003, Eds. Iosif Levin and Gary Gilbert, St. Petersburg, Russia, 2003, 382 pp.

$$R = \frac{2(1-\overline{\mu})^2}{\overline{\mu}(3-\overline{\mu}^2)} \left\{ 1 - \frac{1+\overline{\mu}^2}{\overline{\mu}(3-\overline{\mu}^2)} \ln \left[1 + \frac{\overline{\mu}(3-\overline{\mu}^2)}{1+\overline{\mu}^2} \right] \right\},\tag{4}$$

here $\overline{\mu}$ is defined by Eq. (3).

d) Equation derived in the framework of semi-empirical approach [7] that is based on experimental data presented in Tab. 1:

$$R = \frac{1 - \overline{\mu} / \mu_d}{1 + \overline{\mu} / \mu_u},\tag{5}$$

here

$$\overline{\mu} = a_0 + (1 - a_0) \sqrt{1 - g} + \sum_{n=1}^{6} a_n g^{\frac{n}{3}}, \tag{6}$$

$$\mu_d = \left[1 - \overline{\mu} (1 - \overline{\mu})^2 \sum_{n=0}^{3} b_n \overline{\mu}^{2n} \right] / (2 - \overline{\mu}), \tag{7}$$

$$\mu_{u} = \left[1 - \overline{\mu} \left(1 - \overline{\mu}\right)^{2} \exp\left(\sum_{n=0}^{4} c_{n} \overline{\mu}^{2n}\right)\right] / (2 - \overline{\mu}). \tag{8}$$

The coefficients a_n , b_n and c_n in Eqs. (5-8) are given in Tab. 2.

Table 2. Coefficients to Eqs. (5)-(8).

n	a_n	b_n	C n
0	0.5918	0.0326	-0.0131
1	-0.7937	0.1661	8.4423
2	4.8350	0.7785	-15.6605
3	-22.8150	0.0228	21.8820
4	42.6859		-11.2257
5	-35.8945		
6	11.3905		

Equations (1)-(8) are valid for the full range of variability of parameters g, a, and b_B : $0 \le g \le 1$, $0 \le a \le \infty$, $0 \le b_B \le \infty$. In order to make a comparison and estimate possible errors for the case presented in Fig. 1 we will use also the following equation used in ocean optics research: e) Linear equation by Morel and Prieur [2]:

$$R = g/3. (9)$$

Linear equation (9) was never meant to be used by authors in the full range of variability of g. It was proposed for the case of open ocean waters where g is small.

f) Kubelka-Munk equation [8]:

Proceedings of the II International Conference "Current Problems in Optics of Natural Waters," ONW'2003, Eds. Iosif Levin and Gary Gilbert, St. Petersburg, Russia, 2003, 382 pp.

$$R = \left(1 - \sqrt{1 - g^2}\right) / g. \tag{10}$$

Equation (10) was derived in the assumption that $\overline{\mu}_d = \overline{\mu}_u = 0.5$. Because in seawater both $\overline{\mu}_d$ and $\overline{\mu}_u$ are far from 0.5 (see Tab. 1) that equations is not good for marine optics except special cases with g > 0.6.

g) Equation by Gordon et al. [9] for directed sun illumination:

$$R = 0.0001 + 0.3244 g + 0.1425 g^{2} + 0.1308 g^{3}, (0.1 \le g \le 0.5).$$
 (11)

g) Equation by Gordon et al. [9] for diffuse illumination:

$$R = 0.0003 + 0.3687 g + 0.1802 g^{2} + 0.0740 g^{3}, (0.1 \le g \le 0.5).$$
 (12)

Dependencies of DRC on Gordon's parameter g computed according to Eqs. (1), (2), (4), (5), and (10)-(12) are shown in Fig. 2. Equations (1), (2), (4), and (5) give quite similar results and are valid for the whole range of $0 \le g \le 1$.




Figure 2. Dependence of Diffuse Reflectance Coefficient Gordon's parameter.

4. Possible errors of using linear equations in highly-scattering waters

In order to estimate possible errors to compute diffuse reflection coefficient as a "precise" reference we used values of R as a function of g generated numerically by Hydrolight [10, 11] for the case of diffuse illumination of sea surface. The error distributions for Eqs. (2), (4), (5), and (10)-(12) are shown in Fig. 3. Because the errors of Eq. (1) for the case of diffuse illumination is much smaller, we omitted its dependence from Fig. 3.




Figure 3. Error distribution for various equations to compute diffuse reflectance.

4. Conclusion

The use of linear Eq. (9) is restricted to open ocean waters and coastal waters with $g \le 0.1$. Equation (10) is not adequate for clean ocean waters, and Eqs. (11)-(12) are good only for typical ocean waters excluding very turbid and extremely clean waters [12]. In order to maintain 10% accuracy for the whole range of variability of optical properties of seawater we should use nonlinear Eqs. (1), (2), (4), and (5). The choice of equation should depend on the problem involved because they are related to the different theories.

5. Acknowledgments

The authors thank continuing support at the Naval Research Laboratory (NRL) through the VSF 73-6641-02-5 and HS 73-8028-B2 programs. This article represents NRL contribution PP-7330-03-0010.

6. References

- 1. J. V. R. Zaneveld, "A theoretical derivation of the dependence of the remotely sensed reflectance of the ocean on the inherent optical properties," J. Geophys. Res., 100, 13,135-13,142 (1995).
- 2. A. Morel, L. Prieur, "Analysis of variations in ocean color," *Limnol. Oceanogr.*, 22, 709-722 (1977).
- 3. V. A. Timofeyeva, "Relation Between the Optical Coefficients in Turbid Media," *Izvestiya*, *Atmos. Ocean Physics*, **8**, pp. 654-656, (1972).
- 4. V. A. Timofeyeva, "Determination of Light-Field Parameters in the Depth Regime from Irradiance Measurements," *Izvestiya, Atmos. Ocean Physics*, **15**, 774-776, (1979).
- 5. V. I. Haltrin, "Self-consistent approach to the solution of the light transfer problem for irradiances in marine waters with arbitrary turbidity, depth and surface illumination: I. Case of absorption and elastic scattering," *Appl. Optics*, 37, 3773-3784 (1998).
- 6. V. I. Haltrin, "Exact solution of the characteristic equation for transfer in the anisotropically scattering and absorbing medium." Appl. Optics, 27, 599-602, (1988).
- 7. V. I. Haltrin, "Empirical algorithms to restore a complete set of inherent optical properties of sea water using any two of these properties," *Canadian Journal of Remote Sensing*, **26**, 440-445, (2000).
- 8. P. Kubelka and F. Munk. "Ein Beitrag zur Optik der Farbanstriche," Zeit. Techn. Phys., 12, 593-607 (1931).
- 9. H. R. Gordon, O. B. Brown, and M. M. Jacobs, "Computed relationships between the inherent and apparent optical properties of a flat homogeneous ocean," *Appl. Optics*, **14**, 417-427 (1975).
- 10. C. D. Mobley, Light and Water (Academic Press, San Diego-Toronto, 1994).
- 11. C. D. Mobley, and L. K. Sundman, *Hydrolight 4.1 User Guide*, Sequoia Scientific, Inc., Redmond, WA, 85 pp. (2000).
- 12. The defect in Eqs. (11)-(12) consists of non zero values at g = 0 due to the formal use of regression analysis. The initial unpublished data of Ref. [9] is good for the case of extremely clear water too.